

Quantum Computation of Jones' Polynomials

V. Subramaniam

Department of Electrical Engineering,

*Indian Institute of Technology – Bombay, Mumbai 400076, India**

P. Ramadevi

Department of Physics, Indian Institute of Technology – Bombay, Mumbai 400076, India[†]

(Dated: February 1, 2008)

Abstract

It is a challenging problem to construct an efficient quantum algorithm which can compute the Jones' polynomial for any knot or link obtained from platting or capping of a $2n$ -strand braid. We recapitulate the construction of braid-group representations from vertex models. We present the eigenbases and eigenvalues for the braiding generators and its usefulness in direct evaluation of Jones' polynomial. The calculation suggests that it is possible to associate a series of unitary operators for any braid word. Hence we propose a quantum algorithm using these unitary operators as quantum gates acting on a $2n$ qubit state. We show that the quantum computation gives Jones' polynomial for achiral knots and links.

PACS numbers: 03.67, 03.65

*Electronic address: subbu@ee.iitb.ac.in

[†]Electronic address: ramadevi@phy.iitb.ac.in

I. INTRODUCTION

Quantum algorithms have proven to be more efficient than classical algorithms in solving a number of problems [1]. For instance, Shor's quantum algorithm can factor numbers exponentially faster than classical algorithms. Grover's search algorithm does not change the complexity class, but provides significant speed up for large databases. We would like to explore the power of quantum algorithms in the context of knot theory.

Classification of knots and links in a three-dimensional space is one of the open problems. Jones introduced a recursive procedure for determining a polynomial relation for these knots and links. Jones' polynomials do classify some knots and links [2]. There are other generalized polynomials which improve the classification but none of them have achieved complete classification [3]. It is known that the evaluation of Jones' polynomial classically is a $\#\mathcal{P}$ hard problem [4]. Hence it will be interesting to study the computation of knot and link polynomials using a quantum algorithm.

There are diverse approaches in physics to obtain polynomials for knots and links. Following Alexander's theorem, any knot can be viewed as a closure or capping of an n -strand braid. Therefore the polynomials for knots and links can be determined by studying representation theory of braid groups $\mathcal{B}_n$. The common ingredient in these approaches is to find different representations of braid groups $\mathcal{B}_n$. We now present a brief summary of some of these approaches:

1) **N-state vertex models** which are two-dimensional statistical mechanical models where the bonds of the square lattice carry spin $n/2$ representations of $SU(2)$. The number of possible states of spin $n/2$ is denoted by $N = n + 1$. The properties of these models are described by the so-called R -matrix which is an $N^2 \times N^2$ matrix. The number of nonzero elements in the R -matrix for N -state vertex models is given by $m = \sum_{s=1}^{N-1} 2s^2 + N^2$. In the literature, the vertex models are referred to either as N -state vertex model or m -vertex models; both are equivalent. For example, 2-state vertex models carry spin $1/2$ on the lattice bonds and they are equivalently called as six-vertex models where "six" denotes the number of non-zero R -matrix elements. In Ref. [5], braid group representations and knot polynomials from the R -matrices of $N = 2, 3, 4$ -state vertex models were obtained.

2) **Chern-Simons gauge theory** is a topological field theory which provides a natural framework for the study of knots and links [6]. The knot polynomials are given by the expec-

tation value of Wilson loop observables. In particular, the Jones' polynomial corresponds to the Wilson loop carrying spin $1/2$ representation in $SU(2)$ Chern-Simons theory. Clearly, arbitrary representations of any compact gauge group G can result in generalized polynomials [7]. The polynomials are in the variable q which is a function of the coupling constant k and the rank of the gauge group. The field theoretic polynomials were obtained by exploiting the connection between Chern-Simons theory on a three-manifold with boundary and the corresponding Wess-Zumino-Witten conformal field theory (WZW) on the boundary. The polynomials crucially depended on various representations of the monodromy or braiding matrices in the WZW models.

Recently, Freedman et al. [9] have attempted simulation of topological field theories by quantum computers. The topological quantum computation proposed in Refs. [9]-[11] is at a mathematically abstract level. It exploits the connection between fractional quantum Hall states and Chern-Simons theory at the appropriate integer coupling k .

3) **State sum method** of obtaining bracket polynomials [12, 13]. In Ref. [12], the construction of a unitary representation has been shown for the three-strand braid $\mathcal{B}_3$. Further within this approach, it has been shown that it is not possible for a quantum computer to evaluate the knot polynomial. However, for a specific choice of the polynomial variable, the linking number can be determined [13].

Our aim is to determine the Jones' polynomial for any knot or link obtained from braids using a quantum algorithm. For this purpose, we need to determine matrix representation for braid generators. We will recapitulate the construction of braid group representation from the six-vertex model. Then, we can determine the eigenvalues and eigenstates for these braid generators. This exercise suggests that we can associate product of unitary operators for any braid word. Hence the unitary transformations, corresponding to any braid word, can be implemented using quantum gates. It is important to stress that the quantum computation in this paper is crucially dependent on the mapping of any braid word to a product of unitary operators.

The polynomials for knots and links can be directly computed by choosing a suitable eigenbasis of the braiding matrices. Essentially, a quantum algorithm will determine the probability of finding unitarily evolved initial state ($|in\rangle$) in a final state ($|out\rangle$):

$$P_K = |\langle out | \mathbf{U}_B | in \rangle|^2 \quad (1)$$

where $\mathbf{U}_B$ represents the series of unitary operators corresponding to the braid word. For the braiding matrices obtained from the six-vertex model, the above matrix element gives the modulus-square of the Jones' polynomial (up to an overall constant).

This paper is organized as follows. In Section II, we present a general method to find representation of braid groups using N-state vertex models. We discuss in detail the six-vertex model, braiding eigenvalues and eigenstates. Using these eigenstates, we evaluate Jones' polynomial. In Section III, we present a method to perform the evaluation of the modulus-square of the Jones' polynomial as a quantum computation by considering any knot or link as a composition of cups, a series of braiding operations and caps. In the concluding section, we summarize the results obtained and discuss the significance of the quantum algorithm.

II. N-STATE VERTEX MODEL

In this section, we review the construction of braid group representations from N-state vertex models [5]. In order to compare the eigenstates of the braiding operator with the qubit states, the six-vertex model (spin 1/2 on the bonds of the square lattice) is relevant. Hence, we shall present the explicit form of the R -matrix and the braid matrix for the six-vertex model.

As mentioned in the introduction, vertex models are two-dimensional statistical mechanical models with the spins $n/2$ lying on the bonds of a square lattice. The properties of these models are described by the R -matrix elements between edge states (m_1, m_2) and (n_1, n_2) : $R_{m_1 m_2}^{n_1 n_2}(u)$ where u is the spectral parameter. Here m_1, m_2, n_1, n_2 take values $n/2, n/2 - 1, \dots - n/2$. The integrability condition of these models requires the following equations to be satisfied:

$$X_i(u)X_{i+1}(u+v)X_i(v) = X_{i+1}(v)X_i(u+v)X_{i+1}(u) , \quad (2)$$

$$X_i(u)X_j(v) = X_j(v)X_i(u) \quad |i-j| > 1 , \quad (3)$$

where $X_i(u)$ s are called Yang-Baxter operators and the relation (2) is called Yang-Baxter equation. The explicit form of $X_i(u)$ in terms of the R -matrix elements is given by

$$X_i(u) = \sum_{m_1 m_2 n_1 n_2} R_{m_1 m_2}^{n_1 n_2}(u) I^{(1)} \otimes I^{(2)} \otimes \dots E_{n_2 m_1}^i \otimes E_{n_1 m_2}^{i+1} \otimes \dots I^{(n)} , \quad (4)$$

where $I^{(j)}$ is the identity acting at the j -th position and E is a matrix such that $(E_{n_1 m_2})_{pq} = \delta_{n_1 p} \delta_{m_2 q}$. The solution to Eq. (2) can be written in a compact form [14]:

$$R_{m_1 m_2}^{n_1 n_2}(u) = \sum_{J=0}^n \left\{ \begin{matrix} \frac{n}{2} & \frac{n}{2} & J \\ n_1 & n_2 & m \end{matrix} \right\} \lambda_J(u) \left\{ \begin{matrix} \frac{n}{2} & \frac{n}{2} & J \\ m_2 & m_1 & m \end{matrix} \right\} \quad (5)$$

where the terms in parenthesis are the quantum Clebsch-Gordan coefficients (q-CG) [15] which are nonzero if and only if m takes a value in the range $m = -J, -J + 1, \dots, J$ and satisfies the condition $m_1 + m_2 = m = n_1 + n_2$. Here $\lambda_J(u)$ is given by

$$\lambda_J(u) = \prod_{s=J+1}^{j_1+j_2} \sinh(s\mu + u) \prod_{l=|j_1-j_2|+1}^J \sinh(l\mu - u) , \quad (6)$$

where the q-CG coefficient variable $q = e^{2\mu}$. These solutions are spectral parameter dependent solutions. The explicit form of the R -matrix for the six-vertex model is

$n_1 n_2$	$(\frac{1}{2} \ \frac{1}{2})$	$(\frac{1}{2} \ \frac{-1}{2})$	$(\frac{-1}{2} \ \frac{1}{2})$	$(\frac{-1}{2} \ \frac{-1}{2})$
$m_1 m_2$				
$(\frac{1}{2} \ \frac{1}{2})$	$\sinh(\mu - u)$	0	0	0
$(\frac{1}{2} \ \frac{-1}{2})$	0	$-\sinh(u)$	$e^{-u} \sinh(\mu)$	0
$(\frac{-1}{2} \ \frac{1}{2})$	0	$e^u \sinh(\mu)$	$-\sinh(u)$	0
$(\frac{-1}{2} \ \frac{-1}{2})$	0	0	0	$\sinh(\mu - u)$

We are interested in the construction of braid group representations from these Yang-Baxter operators $X_i(u)$ [5]. The defining relations of the braid group generators are exactly same as Eqs. (2,3) in the absence of the spectral parameter. In other words, the braid group generators $b_i \in \mathcal{B}_n$ (spectral parameter independent operators) are obtained from $X_i(u)$ by taking the spectral parameter $u \rightarrow \infty$:

$$\lim_{u \rightarrow \infty} X_i(u) = b_i . \quad (7)$$

Let us define a quantity $\sigma_{m_1 m_2}^{n_1 n_2}$ as follows:

$$\lim_{u \rightarrow \infty} R_{m_1 m_2}^{n_1 n_2} = \sigma_{m_2 m_1}^{n_1 n_2} . \quad (8)$$

The explicit form of the $\sigma_{m_1, m_2}^{n_1 n_2}$ -matrix elements in this limit turns out to be:

$n_1 n_2$	$(\frac{1}{2} \ \frac{1}{2})$	$(\frac{1}{2} \ \frac{-1}{2})$	$(\frac{-1}{2} \ \frac{1}{2})$	$(\frac{-1}{2} \ \frac{-1}{2})$
$m_1 m_2$				
$(1/2 \ 1/2)$	1	0	0	0
$(1/2 \ -1/2)$	0	0	$-q^{\frac{1}{2}}$	0
$(-1/2 \ 1/2)$	0	$-q^{\frac{1}{2}}$	$1 - q$	0
$(-1/2 \ -1/2)$	0	0	0	1

The above matrix elements indicate that we can choose a basis for the n -strand braid as $|m_1 m_2 \dots m_i m_{i+1} \dots m_n\rangle$ where m_i 's take values $\pm \frac{1}{2}$ which are the two possible quantum states of the spins s_i 's $\equiv \frac{1}{2}$. Hence m_i denotes a single qubit basis and the above basis state is an n -qubit basis state. From Eq. (4), we can write the action of the braiding operator b_i on such an n -qubit state:

$$b_i |m_1 m_2 \dots m_i m_{i+1} \dots m_n\rangle = \sum_{n_i, n_{i+1}} \sigma_{m_i m_{i+1}}^{n_i n_{i+1}} |m_1 m_2 \dots n_i n_{i+1} \dots m_n\rangle, \quad (9)$$

where n_i, n_{i+1} take values $\pm \frac{1}{2}$. Clearly, this equation is not an eigenvalue equation of the braiding operator. We need to diagonalise the $\sigma_{m_1 m_2}^{n_1 n_2}$ matrix to determine the eigenbasis of the braiding operator b_i . As we did in the case of Eq. (5), we can diagonalize the spectral parameter independent $\sigma_{m_1 m_2}^{n_1 n_2}$ using the quantum Clebsch-Gordan coefficients matrix. The corresponding eigenstates will be similar to the coupled states obtained from uncoupled states using quantum CG -coefficient matrix [15]:

$$|J_i m\rangle = \sum_{m_i, m_{i+1}} \begin{Bmatrix} s_i & s_{i+1} & J_i \\ m_i & m_{i+1} & m \end{Bmatrix} |m_i, m_{i+1}\rangle, \quad (10)$$

where s_i, s_{i+1} are the spin $\frac{1}{2}$ states and $J_i \in s_i \otimes s_{i+1} = 0, 1$. Thus we obtain a 2-qubit state $|J_i m\rangle$ which may be an entangled state.

The form of the diagonal σ matrix in the coupled basis is:

$ Jm\rangle$	$ 1 \ 1\rangle$	$ 1 \ 0\rangle$	$ 1 \ -1\rangle$	$ 0 \ 0\rangle$
$ Jm\rangle$				
$ 1 \ 1\rangle$	λ_1	0	0	0
$ 1 \ 0\rangle$	0	λ_1	0	0
$ 1 \ -1\rangle$	0	0	λ_1	0
$ 0 \ 0\rangle$	0	0	0	λ_0

where $\lambda_1 = 1$ and $\lambda_0 = -q$. These eigenvalues are equal up to an overall normalization to the eigenvalues of the Wess-Zumino-Witten model monodromy matrices.

We observe that the eigenvalues of the σ matrix on the coupled states $|Jm\rangle$ depend only on J and not on m . Therefore, we can suppress the m dependence on the eigenstates of the braiding operator and equivalently write it as a tensor product state involving the spin $\frac{1}{2}$ placed on the bonds of the six-vertex model. That is,

$$|J_im\rangle \equiv |J_i\rangle = |(s_i \otimes s_{i+1})_{J_i}\rangle . \quad (11)$$

Even though we have explicitly diagonalized the σ matrix, we must remember that all the braid group generators b_i 's cannot be simultaneously diagonalized. The spectral parameter independent form of Eqs. (2,3) are the defining relations of the braid group $\mathcal{B}_n$ which implies that we can simultaneously diagonalize either b_{2i} 's or b_{2i+1} 's.

A. Representation Theory of Braid Groups in Knot Theory

In this subsection, we would like to address the eigenvectors and eigenvalues of braid generators from the viewpoint of obtaining polynomial invariants of knots from platting or capping of braids.

It is well known that knots from braids are not unique. That is, braids related by Markov moves I and II give rise to the same knot. These two moves indeed completely remove the non-uniqueness. So the construction of polynomial invariants for knots must be such that the polynomial does not change under Markov moves. One such procedure for knots obtained from closure of braids has been presented in [5].

We will use the eigenstates of the braiding operators to directly compute the polynomial invariant for any knot obtained from closure or capping of an n -strand braid. In order to remove the non-uniqueness due to Markov moves, we place orientations on the strands of the braid. Further, we introduce a correction factor to the braid eigenvalues obtained from six-vertex model such that the polynomial does not change under Markov moves I and II. The correction factor on braiding eigenvalues depends on the relative orientations between the two strands. For right-handed half-twists between strands of parallel orientation, the braiding eigenvalues $\lambda_J^{(+)}$ are

$$\lambda_0^{(+)} = -q^{3/2} \ ; \ \lambda_1^{(+)} = q^{1/2} . \quad (12)$$

Similarly, for right-handed half twists between strands of antiparallel orientation, the braiding eigenvalues $\lambda_j^{(-)}$ are

$$\lambda_0^{(-)} = 1 \quad ; \quad \lambda_1^{(-)} = -q^{-1} . \quad (13)$$

The eigenvalues for left-handed half-twists are inverse of the right-handed half-twists eigenvalues.

Suppose we consider any knot obtained from capping of a $2n$ -strand oriented braid. Clearly, capping is possible if the number of outgoing strands is equal the number of incoming strands in the oriented braid. In other words, the quantum states m_i 's on the strands (9) should be such that

$$\sum_{i=1}^{2n} m_i = 0 . \quad (14)$$

Therefore to study knots from braids, only the subspace of the states $|m_1 m_2 \dots m_{2n}\rangle$ satisfying Eq. (14) needs to be considered. Hence the construction of the eigenstates of braiding operators should be consistent with Eq. (14).

We will now present the eigenstates of the braiding matrices which will enable direct evaluation of knot polynomials. For a $2n$ -strand oriented braid, we can write the most general eigenbasis of braiding operators b_{2i+1} with eigenvalue $\lambda_{j_{2i+1}}$, for all i 's, as:

$$\begin{aligned} |\phi_{\{J_{2i+1}\}, \{l_i\}}\rangle = & |(\dots (\dots (s_1 \otimes s_2)_{J_1} \otimes (s_3 \otimes s_4)_{J_3})_{l_1} \\ & \otimes \dots (s_{2i+1} \otimes s_{2i+2})_{J_{2i+1}})_{l_i} \otimes \dots (s_{2n-1} \otimes s_{2n})_{J_{2n-1}})_{l_n} \rangle \end{aligned} \quad (15)$$

Recall that the appropriate braiding eigenvalues (12,13) need to be substituted depending on the relative orientations of the two strands involved in braiding and the handedness. The brackets within the basis kets should be identified with the notation in Eq. (11). That is, $|(s_1 \otimes s_2)_{J_1}\rangle = |J_1\rangle$, $|((s_1 \otimes s_2)_{J_1} \otimes (s_3 \otimes s_4)_{J_3})_{l_1}\rangle = |(J_1 \otimes J_3)_{l_1}\rangle$, and so on. Note that the final combined state in the above basis is chosen to be spin 0 which is essential to satisfy the condition (14) to describe knots from closure or capping of braids.

In the similar fashion, we can write a different eigenbasis for braiding operators b_{2i} with eigenvalue $\lambda_{j_{2i}}$ for all i 's:

$$\begin{aligned} |\tilde{\phi}_{\{J_{2i}\}, \{r_i\}}\rangle = & |((\dots (\dots (s_1 \otimes (s_2 \otimes s_3))_{J_2})_{r_1} \otimes \dots \\ & (s_{2i} \otimes s_{2i+1})_{J_{2i}})_{r_{i-1}} \otimes \dots (s_{2n-2} \otimes s_{2n-1})_{J_{2n-2}})_{r_{n-2}} \otimes s_{2n})_0 \rangle \end{aligned} \quad (16)$$

In order to achieve the final spin 0 state, we require $r_{n-2} = \frac{1}{2}$. Incidentally, these two bases are equivalent to the conformal blocks in Wess-Witten conformal field theory.

The two different bases (15,16) are related by an orthogonal (unitary) duality matrix $a(\{J_{2i+1}\}, \{l_i\}), (\{J_{2i}\}, \{r_i\})$

$$|\phi(\{J_{2i+1}\}, \{l_i\})\rangle = a(\{J_{2i+1}\}, \{l_i\}), (\{J_{2i}\}, \{r_i\}) |\tilde{\phi}(\{J_{2i}\}, \{r_i\})\rangle. \quad (17)$$

The duality matrix can be written in terms of products of $SU(2)_q$ quantum-Racah coefficient matrices [17]:

$$a(\{J_{2i+1}\}, \{l_i\}), (\{J_{2i}\}, \{r_i\}) = \sum_{t_1, \dots, t_{n-2}} \prod_{i=1}^{n-2} \left(a_{t_i J_{2i+1}} \begin{bmatrix} l_{i-1} & s_{2i+1} \\ s_{2i+2} & l_i \end{bmatrix} \prod_{m=0}^{n-2} a_{t_i r_{i-1}} \begin{bmatrix} t_{i-1} & J_{2i} \\ r_i & s_{2m} \end{bmatrix} \right) \prod_{m=0}^{n-2} a_{l_i J_{2i+2}} \begin{bmatrix} t_m & s_{2m+2} \\ s_{2m+3} & t_{m+1} \end{bmatrix} \quad (18)$$

where the closed form expression for the quantum-Racah coefficient matrix is [16]

$$a_{jl} \begin{bmatrix} s_1 & s_2 \\ s_3 & s_4 \end{bmatrix} = (-1)^{s_1+s_2+s_3+s_4} \sqrt{[2j+1][2l+1]} \Delta(s_1, s_2, j) \Delta(s_3, s_4, j) \Delta(s_1, s_4, l) \Delta(s_2, s_3, l) \\ \times \sum_{m \geq 0} (-1)^m [m+1]! \{ [m-s_1-s_2-j]! [m-s_3-s_4-j]! [m-s_1-s_4-l]! \\ \times [m-s_2-s_3-l]! [s_1+s_2+s_3+s_4-m]! [s_1+s_3+j+l-m]! \\ \times [s_2+s_4+j+l-m]! \}^{-1} \quad (19)$$

where $\Delta(a, b, c) = \sqrt{\frac{[-a+b+c]![a-b+c]![a+b-c]!}{[a+b+c+1]!}}$ and the number in square brackets represents the q -number defined as

$$[x] = \frac{q^{x/2} - q^{-x/2}}{q^{1/2} - q^{-1/2}}.$$

With this detailed background on the representation theory of braid groups, we are now in a position to evaluate Jones' polynomial for any knot or link.

B. Evaluation of Jones' polynomial

Consider any knot or link as shown in Fig. 1 which is technically called platting or capping of braids. The knot type is determined by the braiding region denoted as the shaded region in Fig. 1. It involves a sequence of braiding operations which is usually written as a braid




FIG. 1: Viewing Knots as a Capping of Braiding Sequences

word with appropriate orientations. Taking the initial state and final state as $|\phi_{(\{0\},\{0\})}\rangle$, the following matrix element

$$V_K = \langle \phi_{(\{0\},\{0\})} | \mathbf{B} | \phi_{(\{0\},\{0\})} \rangle \quad (20)$$

gives the Jones' polynomial (up to an overall normalisation). To see the polynomial form in variable q , we can substitute the eigenvalues of the braiding generators and the unitary duality matrices elaborated in the previous subsection.

For the variable q being root of unity, we can equivalently denote the sequence of braiding operations $\mathbf{B} \in \mathcal{B}_{2n}$ as a product of unitary $2^{2n} \times 2^{2n}$ matrices. We will now illustrate this mapping for two examples involving four-strand and six-strand braids. As shown in Fig.2(a), the braid word

$$B_a = b_2^3 \hat{b}_1^{-2} \hat{b}_3^{-2} b_2^3 \in \mathcal{B}_4 ,$$

where $\hat{b}_i$ denotes braiding between anti-parallelly oriented i -th strand and $i+1$ -th strand and b_j denotes braiding between parallelly oriented j -th strand and $j+1$ -th strand. Following the rules of the representation theory of braids, the braid-word in Fig. 2(a) will be mapped to a product of $2^4 \times 2^4$ unitary matrices as follows:

$$B_a \equiv \mathbf{U}_{B_a} = (\{\mathbf{a} \mathbf{f}(\mathbf{b}_2) \mathbf{a}^\dagger \mathbf{g}(\mathbf{b}_1, \mathbf{b}_3) \mathbf{a} \mathbf{h}(\mathbf{b}_2) \mathbf{a}^\dagger\}_{2 \times 2} \oplus \mathbf{1}_{14 \times 14})_{2^4 \times 2^4} \quad (21)$$

where $\mathbf{a}$ is the duality matrix and the matrix elements of the matrices $\mathbf{f}(\mathbf{b}_2)$, $\mathbf{g}(\mathbf{b}_1, \mathbf{b}_3)$ and




FIG. 2: Two examples (a) $B_a \in \mathcal{B}_4$, (b) $B_b \in \mathcal{B}_6$

$\mathbf{h}(\mathbf{b}_2)$, for the braid word B_a , will be

$$\mathbf{f}(\mathbf{b}_2)_{J_1 J_2} = \delta_{J_1 J_2} (\lambda_{J_2}^{(+)})^3 , \quad \mathbf{g}(\mathbf{b}_1, \mathbf{b}_3)_{J_1, J_2} = \delta_{J_1 J_2} (\lambda_{J_1}^{(-)})^{-4} , \quad \mathbf{h}(\mathbf{b}_2)_{J_1 J_2} = \delta_{J_1 J_2} (\lambda_{J_2}^{(+)})^3 . \quad (22)$$

The entries in these diagonal matrices are functions of the braiding eigenvalues. The relative orientations and the number of crossings between the respective braids are incorporated in writing the functional form. If q is a root of unity, these diagonal matrices are unitary. Thus we see that the braid word $B_a \in \mathcal{B}_4$ can be equivalently represented as a product of unitary $2^4 \times 2^4$ matrices. Similarly, the braid word B_b corresponding to Fig. 2(b) will be

$$B_b = (b_2^{-1} b_4) (\hat{b}_1^{-2} \hat{b}_3^{-3} \hat{b}_5^{-2}) (\hat{b}_2 \hat{b}_4^2) b_1 \hat{b}_2 \in \mathcal{B}_6 .$$

The equivalent description involving product of $2^6 \times 2^6$ unitary matrices will be

$$\mathbf{U}_{B_b} = [(\mathbf{a} \mathbf{f}(\mathbf{b}_2, \mathbf{b}_4) \mathbf{a}^\dagger \mathbf{g}(\mathbf{b}_1, \mathbf{b}_3, \mathbf{b}_5) \mathbf{a} \mathbf{h}(\mathbf{b}_2, \mathbf{b}_4) \mathbf{a}^\dagger \mathbf{f}_1(\mathbf{b}_1) \mathbf{a} \mathbf{g}_1(\mathbf{b}_2) \mathbf{a}^\dagger)_{5 \times 5} \oplus \mathbf{1}_{59 \times 59}]_{2^6 \times 2^6} \quad (23)$$

where the duality matrix elements are denoted as $a_{(J_1, J_3, J_5=l_1), (J_2, J_4, r_1)}$ and the remaining 5×5 diagonal matrices for the braid word B_b will be

$$\begin{aligned}
\mathbf{f}(\mathbf{b}_2, \mathbf{b}_4)_{(J'_2, J'_4, r'_1), (J_2, J_4, r_1)} &= \delta_{J'_2 J_2} \delta_{J'_4 J_4} \delta_{r'_1, r_1} (\lambda_{J_2}^{(+)})^{-1} \lambda_{J_4}^{(+)} \\
\mathbf{g}(\mathbf{b}_1, \mathbf{b}_3, \mathbf{b}_5)_{(J'_1, J'_3, J'_5), (J_1, J_3, J_5)} &= \delta_{J'_1 J_1} \delta_{J'_3 J_3} \delta_{J'_5 J_5} (\lambda_{J_1}^{(-)})^{-2} (\lambda_{J_3}^{(-)})^{-3} (\lambda_{J_5}^{(-)})^{-2} \\
\mathbf{h}(\mathbf{b}_2, \mathbf{b}_4)_{(J'_2, J'_4, r'_1), (J_2, J_4, r_1)} &= \delta_{J'_2 J_2} \delta_{J'_4 J_4} \delta_{r'_1, r_1} \lambda_{J_2}^{(-)} (\lambda_{J_4}^{(-)})^2 \\
\mathbf{f}_1(\mathbf{b}_1)_{(J'_1, J'_3, J'_5), (J_1, J_3, J_5)} &= \delta_{J'_1 J_1} \delta_{J'_3 J_3} \delta_{J'_5 J_5} \lambda_{J'_1}^{(+)} \\
\mathbf{g}_1(\mathbf{b}_2)_{(J'_2, J'_4, r'_1), (J_2, J_4, r_1)} &= \delta_{J'_2 J_2} \delta_{J'_4 J_4} \delta_{r'_1, r_1} \lambda_{J'_2}^{(-)} .
\end{aligned} \tag{24}$$

In a similar fashion, we can find the unitary representation $\mathbf{U}_B$ for any braid word $B \in \mathcal{B}_{2n}$ in terms of products of $2^{2n} \times 2^{2n}$ unitary matrices involving duality matrices $[a_{(\{J_{2i+1}\}, \{l_i\}), (\{J_{2i}\}, \{r_i\})} \oplus \mathbf{1}]_{2^{2n} \times 2^{2n}}$ and diagonal braiding matrices. These unitary representations play the role of quantum gates in the quantum computation of Jones' polynomial.

III. QUANTUM COMPUTATION

In this section, we attempt to compute the Jones' polynomials, for knots and links obtained from platting or capping of $2n$ strand braid as shown in Fig. 1, through a quantum algorithm. We have already elaborated in the previous section that we can associate $\mathbf{U}_B$ (product of unitary matrices) for every braid word $B \in \mathcal{B}_{2n}$. The quantum algorithm involves the following steps:

Step 1: Let the initial $2n$ -qubit state be $|0\rangle$ ($|cup\rangle$).

Step 2: We perform the sequence of unitary operations $\mathbf{U}_B$ corresponding to the braid word $B \in \mathcal{B}_{2n}$ in Fig. 1. The unitarily transformed state will be

$$|\Psi\rangle = \mathbf{U}_B |0\rangle . \tag{25}$$

Step 3: Finally, we determine the probability of the unitarily evolved state $|\Psi\rangle$ in a specific final state $|f\rangle$ as

$$|\langle f | \mathbf{U}_B | 0 \rangle|^2 . \tag{26}$$

Taking the final state to be $|cap\rangle = |0\rangle$, we get the modulus square of the Jones' polynomial (up to an overall normalisation) V_K (20).

For a subclass of knots(links) called achiral knots(links), V_K is unchanged under $q \rightarrow q^{-1}$. In other words, the matrix element $\langle 0|U_B|0\rangle$ will be real. For these achiral knots and links, the quantum algorithm directly gives the Jones' polynomial (up to an overall normalisation).

IV. SUMMARY AND DISCUSSIONS

In this paper, we have presented matrix representations for braiding matrices from six-vertex models. We have discussed the representation theory of braids, namely, the eigenbasis and eigenvalues of the braid generators obtained from six-vertex models. The explicit evaluation of Jones' polynomial, for any knot/link from braids, is presented. From the evaluation, we have shown that we can associate a series of unitary operators for any braid word. This is the significant result of the paper enabling quantum computation. We have demonstrated a quantum algorithm, involving these unitary operators, which can determine the modulus square of the Jones' polynomial for any knot or link. The algorithm gives Jones' polynomial for achiral knots and links.

We must realize that the quantum computation essentially determines the probability of unitarily evolved initial state in a specific final state. Further, the number of unitary operators is dependent on the braid word and at most equal to twice the length of the braid word.

Acknowledgments: We would like to thank L.H. Kauffman for comments and queries which significantly helped us to improve the paper. We would also like to thank Umasankar for going over the manuscript and suggesting corrections.

-
- [1] A. Ekert, P. Hayden, and H. Inamori, quant-ph/0011013 and references therein.
A. Ekert and R. Jozsa, Reviews of Modern Physics **68** (1996) 733-753.
 - [2] V.F.R. Jones, Bull. AMS **12** (1985) 103-111.
 - [3] P. Freyd, D. Yetter, J. Hoste, W.B.R. Lickorish, K. Millet and A. Ocaneu, Bull. AMS **12** (1985) 239-246.
L.H. Kauffman, Ann. Math. Studies (Princeton Univ. Press, 1987); (World Scientific, 1991).
 - [4] F. Jaeger, D. Vertigen, and D. Welsh, Math. Proc. Cambridge Philos. Soc. **108** (1990)35-53.

- [5] Y. Akutsu and M. Wadati, Jour. of the Phys. Soc. of Japan, **56** (1987) 3039-3051; M. Wadati, T. Deguchi and Y. Akutsu, Phys. Rept. **180** (1989) 247-332.
- [6] E. Witten, Commun. Math. Phys. **121** (1989) 351-391.
- [7] P. Ramadevi, T.R. Govindarajan and R.K. Kaul, Nucl. Phys. B **402** (1993) 548-566.
- [12] L. H. Kauffman, math.QA/0105255.
- [9] M.H. Freedman, A. Kitaev, and Z. Wang, Commun. Math. Phys. **227** (2002) 587-603, quant-ph/0001071.
- [10] M.H. Freedman, quant-ph/0003128.
- [11] M.H. Freedman, A. Kitaev, M.J. Larsen, Z. Wang, quant-ph/0101025.
- [12] L. H. Kauffman, math.QA/0105255.
- [13] L. H. Kauffman and S. J. Lomonaco Jr., quant-ph/0205137.
- [14] R.K. Kaul, hep-th/9804122, Frontiers of Field Theory, Quantum Gravity and Strings, Nova Science, (1999) 45; P. Ramadevi, Ph.D Thesis (1996).
- [15] V. Pasquier, Commun. Math. Phys. **118** (1988) 355-364.
- [16] A.N. Kirillov, N.Yu. Reshetikhin, ed. V.G. Kac, World Scientific, 1989.
- [17] R.K. Kaul, Commun. Math. Phys. **162** (1994) 289-320.
- [18] L.M.K. Vandersypen, M. Steffen, G. Breyta, C.S. Yannoni, M.H. Sherwood, and I.L. Chaung, Nature **414** (2001) 883-887.